

2A: Solving Quadratic Equations

When solving quadratic equations, we have several methods that we can use depending on the terms of the quadratic expression. Below is a summary of these methods with an example.

Standard Form of a quadratic equation: $ax^2 + bx + c = 0$ for integers a, b, c .

Method 1: Square Root Method

$$ax^2 + c = 0$$

Example: Solve $4x^2 - 27 = 0$

$$4x^2 - 27 = 0$$

$$4x^2 = 27$$

$$x^2 = \frac{27}{4}$$

Isolate the x^2

$$x = \pm \sqrt{\frac{27}{4}}$$

Square Root

$$x = \pm \frac{3\sqrt{3}}{2}$$

Simplify

Method 2: Factoring

The factoring method employs the zero-product rule which states:

“If $pq = 0$, then either $p = 0$, or $q = 0$.”

When $a = 1$: $x^2 + bx + c = 0$

Strategy: Find two numbers m and n such that

$$m + n = b, \quad \text{and} \quad m \cdot n = c$$

then we write

$$x^2 + bx + c = (x + m)(x + n) = 0$$

and use the zero product rule.

Example: Solve $x^2 - 2x - 7 = 8$

$$x^2 - 2x - 7 = 8$$

$$x^2 - 2x - 15 = 0$$

Make one side equal to 0

$$(x - 5)(x + 3) = 0$$

Factor the quadratic expression

$$x - 5 = 0, \quad \text{or} \quad x + 3 = 0$$

Set each factor equal to zero.

$$x = 5, \quad \text{or} \quad x = -3$$

Solve each sub-equation to get solutions.

When $\neq 1$, use the "ac" method: $ax^2 + bx + c = 0$

Example: Solve $2x^2 + x - 15 = 0$

$$2x^2 + x - 15 = 0$$

First make sure that the equation is equal to 0.

Compute the value of ac

$$ac = 2(-15) = -30$$

Find two numbers that have a product of $ac = -30$ and a sum of $b = 1$. They are -5 and 6.

$$2x^2 + 6x - 5x - 15 = 0$$

Split the bx term into a sum with these coefficients.

$$2x(x + 3) - 5(x + 3) = 0$$

Factor the expression by *grouping*.

$$(2x - 5)(x + 3) = 0$$

$$2x - 5 = 0, \text{ or } x + 3 = 0$$

Set both factors equal to zero.

$$x = \frac{5}{2}, \text{ or } x = -3$$

Solve the sub-equations.

Square Trinomials: $a^2x^2 + 2abx + b^2 = c$

Example: Solve $9x^2 + 30x + 25 = 40$

$$9x^2 + 30x + 25 = 40$$

First check to see if the left side is a quadratic trinomial in the form $a^2x^2 + 2abx + b^2 \dots$ *This one is!*

$$(3x + 5)^2 = 40$$

Factor the expression.

$$3x + 5 = \pm\sqrt{40}$$

Square root (don't forget the \pm roots)

$$3x = \pm\sqrt{40} - 5$$

Isolate the x variable to solve.

$$x = \frac{\pm 2\sqrt{10} - 5}{3}$$

Method 3: Completing the Square

Use for any quadratic equation in standard form $ax^2 + bx + c = 0$.

Example: Solve $3x^2 + 24x - 21 = 0$

$$3x^2 + 24x - 21 = 0$$

$$3x^2 + 24x = 21$$

Isolate constant term c

$$x^2 + 8x = 7$$

Divide by $a = 3$

$$x^2 + 8x + 16 = 23$$

Add $(b/2)^2$ to both sides, where b is the coefficient on x

$$(x + 4)^2 = 23$$

Factor the perfect square trinomial

$$x + 4 = \pm\sqrt{23}$$

Square root both sides

$$x = \pm\sqrt{23} - 4$$

Subtract 4 to solve.

Method 4: The Quadratic Formula

Use for any quadratic in standard form $ax^2 + bx + c = 0$.

The solutions to $ax^2 + bx + c = 0$ can be found using the formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Example: Solve $2x^2 + 4x - 5 = 6$

$$2x^2 + 4x - 5 = 6$$

$$2x^2 + 4x - 11 = 0$$

Set equation equal to 0

$$x = \frac{-(4) \pm \sqrt{(4)^2 - 4(2)(-11)}}{2(2)}.$$

Use coefficients $a = 2, b = 4, c = -11$ for the formula.

$$x = \frac{-4 \pm \sqrt{16 + 88}}{4}$$

$$x = \frac{-4 \pm \sqrt{104}}{4}$$

Simplify *carefully!*

$$x = \frac{-4 \pm 2\sqrt{26}}{4}$$

$$x = \frac{-2 \pm \sqrt{26}}{2}$$

Complex Solutions

In the set of real numbers the square root of all real numbers is a real number. However, the square root of a negative real number is undefined. This is because when we square any real number, we get a non-negative number.

So, we have to define a new type of number called an **imaginary number**. So, we define

$$\sqrt{-1} = i$$

to be our imaginary unit. An **imaginary number** is any multiple of i .

For example:

Write the $\sqrt{-4}$ and $\sqrt{-8}$ using the imaginary unit i .

$$\sqrt{-4} = \sqrt{4} \cdot \sqrt{-1} = 2i$$

$$\sqrt{-8} = \sqrt{4} \cdot \sqrt{-1} \cdot \sqrt{2} = 2 \cdot i \cdot \sqrt{2} = 2i\sqrt{2}$$

A **complex number** has two parts: a real part and an imaginary part of the form

$$\underbrace{a}_{\text{real part}} + \underbrace{bi}_{\text{imag. part}}$$

When solving quadratic equations, we sometimes end up with no real solutions but we have imaginary solutions.

Example. Use the square root method to solve $4x^2 + 20 = 10$

$$4x^2 + 20 = 10$$

$$x^2 = \frac{-10}{4}$$

$$x = \sqrt{\frac{-10}{4}} = \frac{\sqrt{-10}}{2} = \frac{\sqrt{10}}{2} \cdot \sqrt{-1} = \frac{\sqrt{10}}{2}i$$

Assignment 2A

Solve using the square root method

1. $3(x + 3)^2 - 5 = 31$

2. $2x^2 + 9 = 5$

3. $(x - 4)^2 = -9$

Solve by Factoring

1. $x^2 + 15x - 34 = 0$

2. $x^2 - 12x = -35$

3. $4x^2 + 12x + 9 = 0$

4. $2x^2 + 5x + 3 = 0$

5. Try this: Factor the left side *before* setting equal to 0.

$9x^2 - 24x + 16 = -5$

Solve by completing the square

6. $2x^2 + 8x - 5 = 7$

7. $x^2 + 6x = -25$

8. $4x^2 + 8x + 20 = 0$

Solve using the quadratic formula

9. $4x^2 - 3x - 5 = 2$

10. $5x^2 - 4x = 5$

11. $3x^2 + x + 1 = 0$

12. $-5x^2 + 6x + 3 = 5$