

# 1E: Difference Quotient

## Notes

A very important question in mathematics is how to find the slope of a line that is tangent to a curve. A tangent line touches a curve in exactly one point without crossing the curve. In formal Geometry courses we study lines that are tangent to a circle and discover some interesting results such as the fact that a tangent line to a circle is perpendicular to the radius that intersects it.

In the previous lesson, we found the average rate of change. Graphically, the rate of change on the interval  $[a, b]$  represents the slope of the secant line through the points  $a$  and  $b$  (as in the example to the right)


We now want to choose two points that are closer and closer together. The example to the right shows a secant line for the interval  $[3, 5]$ . We may want to also consider the intervals

$$[3, 4], [3, 3.5], [3, 3.1], \dots, [3, 3.000001].$$

What would happen to the slope of the secant line of

$$f(x) = (x - 3)^2$$

over these intervals listed above? (Use the graph on the right to help.)



To generalize this, we will set the distance between the two endpoints and call this  $h$ . This leaves us with the interval  $[x, x + h]$ . Now, we will find is the rate of change for  $f(x)$  on the interval  $[x, x + h]$ . We will call this function  $D(x)$  the **Difference Quotient**

$$D(x) = \frac{\Delta y}{\Delta x} = \frac{f(x + h) - f(x)}{(x + h) - x}$$

or

**Difference Quotient:**

$$D(x) = \frac{f(x + h) - f(x)}{h}$$

Example:

Find and simplify the difference quotient function  $D(x)$  for each function.

a)  $f(x) = 2x + 3$

b)  $g(x) = x^2$

c) Use the result of (b) to find the rate of change for  $g(x) = x^2$  on  $[1,3]$ .

**Limit as  $h \rightarrow 0$**

In the difference quotient function,  $h$  cannot be 0 since it is in the denominator. However, an important question to ask is

*"What does the value of the function become as  $h$  goes to 0?"*

Example:

Consider the function  $D(x)$  above for  $g(x) = x^2$ .

What does this function become as  $h$  goes to zero?

**\*\*Key Point:** The value of  $D(a)$  as  $h \rightarrow 0$  is the \_\_\_\_\_  
of the line *tangent* to the curve at  $(a, f(a))$ .

### Assignment:

For each of the functions below, find and simplify  $D(x)$ .

1.  $a(x) = 3x - 2$

2.  $b(x) = \frac{1}{2}x + 4$

3.  $c(x) = mx + b$ . (treat  $m$  and  $b$  as numbers)

4.  $h(x) = 2x^2$

5.  $j(x) = 3x^2$

6.  $k(x) = x^2 + x$

7.  $m(x) = x^2 + 5$